

Coding Resource

INDICATION

ETCAMAH in combination with a CDK4/6 inhibitor (abemaciclib, palbociclib, or ribociclib) is indicated for the treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer upon detection of *ESR1* mutation during aromatase inhibitor and CDK4/6 inhibitor therapy, based on an FDA-authorized test.

This indication is approved under accelerated approval based on progression-free survival as measured from detection of *ESR1* mutation. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

It is important to note that the codes identified below are examples only. Each provider is responsible for ensuring all coding is accurate and documented in the medical record based on the condition of the patient. The use of the following codes does not guarantee reimbursement.

National Drug Code (NDC)¹

10-digit NDC

Dosage	Code
75 mg tablets	0310-0075-01

11-digit NDC

Dosage	Code
75 mg tablets	00310-0075-01

Diagnosis Codes²

ICD-10-CM	Description
PRIMARY BREAST CANCER SITE	
C50.011	Malignant neoplasm of nipple and areola, right female breast
C50.012	Malignant neoplasm of nipple and areola, left female breast
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast
C50.021	Malignant neoplasm of nipple and areola, right male breast
C50.022	Malignant neoplasm of nipple and areola, left male breast
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast
C50.111	Malignant neoplasm of central portion of right female breast
C50.112	Malignant neoplasm of central portion of left female breast
C50.119	Malignant neoplasm of central portion of unspecified female breast
C50.121	Malignant neoplasm of central portion of right male breast
C50.122	Malignant neoplasm of central portion of left male breast
C50.129	Malignant neoplasm of central portion of unspecified male breast
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast

ICD-10-CM=International Classification of Diseases, Tenth Revision, Clinical Modification.

Please see Important Safety Information on pages 4-6 and complete Prescribing Information, including Boxed WARNING, and Patient Information.

Coding Resource

Diagnosis Codes² (cont'd)

ICD-10-CM	Description
PRIMARY BREAST CANCER SITE (cont'd)	
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast
C50.611	Malignant neoplasm of axillary tail of right female breast
C50.612	Malignant neoplasm of axillary tail of left female breast
C50.619	Malignant neoplasm of axillary tail of unspecified female breast
C50.621	Malignant neoplasm of axillary tail of right male breast
C50.622	Malignant neoplasm of axillary tail of left male breast
C50.629	Malignant neoplasm of axillary tail of unspecified male breast
C50.811	Malignant neoplasm of overlapping sites of right female breast
C50.812	Malignant neoplasm of overlapping sites of left female breast
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast
C50.821	Malignant neoplasm of overlapping sites of right male breast
C50.822	Malignant neoplasm of overlapping sites of left male breast

Please see Important Safety Information on pages 4-6 and complete Prescribing Information, including **Boxed WARNING, and **Patient Information**.**

Coding Resource

Diagnosis Codes² (cont'd)

ICD-10-CM	Description
PRIMARY BREAST CANCER SITE (cont'd)	
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast
C50.911	Malignant neoplasm of unspecified site of right female breast
C50.912	Malignant neoplasm of unspecified site of left female breast
C50.919	Malignant neoplasm of unspecified site of unspecified female breast
C50.921	Malignant neoplasm of unspecified site of right male breast
C50.922	Malignant neoplasm of unspecified site of left male breast
C50.929	Malignant neoplasm of unspecified site of unspecified male breast
PERSONAL HISTORY OF MALIGNANT NEOPLASM OF BREAST	
Z85.3	Personal history of malignant neoplasm of breast
ESTROGEN AND OTHER HORMONE & GROWTH FACTOR RECEPTORS	
Z17.0	Estrogen receptor positive status [ER+]
Z17.32	Human epidermal growth factor receptor 2 negative status
Z17.411	Hormone receptor positive with human epidermal growth factor receptor 2 negative status
METASTASIS TO BONE AND BONE MARROW	
C79.51	Secondary malignant neoplasm of bone
C79.52	Secondary malignant neoplasm of bone marrow
METASTASIS TO LUNG AND PLEURA	
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.1	Secondary malignant neoplasm of mediastinum
C78.2	Secondary malignant neoplasm of pleura
C78.30	Secondary malignant neoplasm of unspecified respiratory organ
C78.39	Secondary malignant neoplasm of other respiratory organs
METASTASIS TO BRAIN	
C79.31	Secondary malignant neoplasm of brain
C79.32	Secondary malignant neoplasm of cerebral meninges
METASTASIS TO BRAIN	
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct

Please see Important Safety Information on pages 4-6 and complete **Prescribing Information**, including **Boxed WARNING**, and **Patient Information**.

IMPORTANT SAFETY INFORMATION INCLUDING BOXED WARNING ABOUT ETCAMAH® (camizestrant)

WARNING: ARRHYTHMIA RISK WITH CONCOMITANT USE OF QTc INTERVAL PROLONGING DRUGS

ETCAMAH in combination with ribociclib, a QTc interval prolonging drug and a strong CYP3A inhibitor, or in combination with other QTc interval prolonging drugs, can increase the risk of Torsades de Pointes (TdP), other ventricular arrhythmias, and sudden death.

Avoid concomitant use of ETCAMAH with products (other than ribociclib) known to prolong the QTc interval and/or have a known risk of TdP. If concomitant use with other products cannot be avoided, monitor the QTc interval more frequently.

Obtain electrocardiogram (ECG) prior to initiation and monitor heart rate (HR) and QTc interval during treatment. Assess and correct electrolyte abnormalities prior to initiation and during treatment. Withhold ETCAMAH until resolution of QTc interval prolongation and resume or permanently discontinue ETCAMAH based on severity.

WARNINGS AND PRECAUTIONS

QTc Interval Prolongation: ETCAMAH in combination with a CDK4/6 inhibitor (CDK4/6i) is associated with QTc interval prolongation. When ETCAMAH is used in combination with ribociclib, a CDK4/6i, there is potential for increased risk of TdP, other ventricular arrhythmias, and sudden death. One Grade 4 case of TdP was observed in a dose-finding trial when ETCAMAH was used with ribociclib. In SERENA-6, QTc interval prolongation occurred in 2.6% of patients treated with ETCAMAH in combination with a CDK4/6i. QTc interval prolongation led to dose interruption in 0.6% of patients. No patients in SERENA-6 discontinued ETCAMAH due to QTc interval prolongation. ETCAMAH also causes bradycardia, which increases the risk of QTc interval prolongation.

Perform an ECG prior to initiating ETCAMAH, then every week for the first 2 weeks of treatment, and periodically during treatment as clinically indicated. Obtain serum electrolytes at baseline and during treatment as clinically indicated, and correct electrolyte abnormalities. Avoid concomitant use of ETCAMAH in combination with a CDK4/6i with products that cause QTc interval prolongation, are strong CYP3A inhibitors, and/or are drugs known to lower HR.

For QTc >500 msec or QTc >480 msec and prolongation from baseline >60 msec, withhold ETCAMAH. If other contributing causes are identified, then treat or correct the contributing causes. Resume ETCAMAH when QTc returns to <480 msec. Reassess ECGs weekly for the first 2 weeks of treatment, and periodically during treatment as clinically indicated. Permanently discontinue ETCAMAH if QTc interval prolongation is either >500 msec or >60 msec change from baseline AND associated with any: TdP, polymorphic ventricular tachycardia, syncope, or signs/symptoms of serious arrhythmia.

Bradycardia: ETCAMAH causes a decrease in HR. Bradycardia increases the risk for life-threatening arrhythmias and sudden death when concomitant QTc interval prolongation is present, such as when ETCAMAH is used in combination with ribociclib, both a QTc interval prolonging product and a strong CYP3A inhibitor. In SERENA-6, bradycardia occurred in 8% of patients. The mean HR decrease from baseline was approximately 13 beats per minute (bpm) with the maximum decrease observed on day 15. Median time to onset was 17 days (range 13 to 283) after starting ETCAMAH. No patients discontinued ETCAMAH due to bradycardia. Dose interruption occurred in 3.9% of patients. The safety of ETCAMAH has not been established in patients with a baseline resting HR <55 bpm as these patients were excluded from SERENA-6. Monitor HR more frequently during the first 30 days of treatment in patients with bradycardia (HR <60 bpm) at baseline and those on concomitant medications known to lower HR (eg, beta-blockers).

For symptomatic (Grade 2 or above) bradycardia, withhold ETCAMAH until symptoms resolve. Obtain ECG to evaluate etiology. If a contributing concomitant medication is identified, modify the dosage or discontinue this medication, as appropriate, until bradycardia symptoms resolve, then resume ETCAMAH. Consider reassessing the HR after restart. Permanently discontinue for persistent symptomatic bradycardia.

Embryo-Fetal Toxicity: Based on findings in animals and its mechanism of action, ETCAMAH can cause fetal harm when administered to a pregnant woman. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective non-hormonal contraception during treatment with ETCAMAH and for 4 weeks after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ETCAMAH and for 1 week after the last dose.

IMPORTANT SAFETY INFORMATION INCLUDING BOXED WARNING ABOUT ETCAMAH® (camizestrant) (cont'd)

ADVERSE REACTIONS

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, with ETCAMAH in combination with a CDK4/6i were decreased neutrophils (68%), decreased leukocytes (66%), decreased hemoglobin (47%), decreased lymphocytes (39%), decreased platelets (36%), visual disturbances (34%), and fatigue (23%).

Permanent discontinuation due to adverse reactions occurred in 1.3% of patients. Adverse reactions that resulted in permanent discontinuation of ETCAMAH included gastroesophageal reflux disease, cholestasis and hepatic cytolysis (0.6% each). Dosage interruption of ETCAMAH due to adverse reactions occurred in 22% of patients.

Visual disturbances: For visual disturbances limiting instrumental ADLs (activities of daily living) (Grade 2) or above, withhold ETCAMAH until symptoms resolve to Grade 1 or below. Refer to an eye professional for an ophthalmic examination and treatment and reassess at the next visit.

For other Grade 3 or higher adverse reactions: Withhold ETCAMAH until resolution to Grade 2 or below, then resume. Permanently discontinue for recurrence of Grade 3 or higher adverse reactions.

DRUG INTERACTIONS

- **Strong CYP3A Inhibitors:** Monitor for increased adverse reactions to ETCAMAH and modify the dosage as recommended
- **Strong and Moderate CYP3A Inducers:** Avoid the use of strong CYP3A inducers. Use caution with the co-administration of a moderate CYP3A inducer
 - Avoid concomitant use of moderate CYP3A inducers for patients who are receiving ETCAMAH in combination with abemaciclib or palbociclib. If concomitant use cannot be avoided, increase the ETCAMAH dosage from 75 mg once daily to 150 mg once daily. After the moderate CYP3A inducer has been discontinued for at least 14 days, resume the ETCAMAH dosage used prior to initiation of the moderate CYP3A inducer
 - For patients who are receiving ETCAMAH in combination with ribociclib concomitantly with a moderate CYP3A inducer, no ETCAMAH dosage modification is recommended
- **CYP2C9 and/or CYP2C19 Substrates:** Avoid concomitant use of ETCAMAH with CYP2C9 or CYP2C19 substrates during treatment with ETCAMAH and at least 2 weeks after the last dose of ETCAMAH, unless otherwise recommended in the Prescribing Information of the CYP2C9 or CYP2C19 substrate
- **Certain CYP3A Substrates:** Refer to the Prescribing Information for CYP3A substrates where minimal increases in the concentration may lead to serious adverse reactions
- **Drugs that Prolong the QTc Interval:** ETCAMAH is indicated in combination with a CDK4/6i and there is increased risk of QTc interval prolongation with ribociclib, a strong CYP3A inhibitor that can prolong the QTc interval. Refer to the ribociclib Prescribing Information for dosage modifications. Avoid concomitant use of ETCAMAH with products (other than ribociclib) known to prolong the QTc interval and/or have a known risk of TdP. If concomitant use with other products cannot be avoided, monitor the QTc interval more frequently. ETCAMAH in combination with a CDK4/6i is associated with QTc interval prolongation
- **Drugs that Cause Bradycardia:** Avoid concomitant use of ETCAMAH with other products known to cause bradycardia. Monitor for signs and symptoms of bradycardia if concomitant use cannot be avoided. ETCAMAH causes decreases in HR that are dose and baseline HR dependent

Refer to the Prescribing Information for the co-administered CDK4/6i for dosage modification guidelines related to adverse reactions, organ impairment, and drug-drug interactions.

SPECIAL POPULATIONS

Pregnancy: Based on findings in animals and mechanism of action, ETCAMAH can cause fetal harm when administered to a pregnant woman. Advise pregnant women and females of reproductive potential of the potential risk to a fetus.

Lactation: Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ETCAMAH and for 1 week after the last dose.

Please see full **Prescribing Information**, including **Boxed WARNING**, and **Patient Information**.

IMPORTANT SAFETY INFORMATION INCLUDING BOXED WARNING ABOUT ETCAMAH® (camizestrant) (cont'd)

SPECIAL POPULATIONS (cont'd)

Females and Males of Reproductive Potential: ETCAMAH can cause fetal harm when administered to pregnant women. Verify pregnancy status of female patients of reproductive potential prior to initiating ETCAMAH. Advise female patients of reproductive potential to use effective non-hormonal contraception during treatment with ETCAMAH and for 4 weeks after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ETCAMAH and for 1 week after the last dose. Refer to the Prescribing Information of the CDK4/6i palbociclib, if used in combination with ETCAMAH, for contraception information and use for the longest recommended post-treatment duration. ETCAMAH may impair fertility in female and male patients.

Pediatric Use: Safety and effectiveness have not been established in pediatric patients.

Hepatic Impairment: In patients with severe (Child-Pugh C) hepatic impairment, when ETCAMAH is combined with ribociclib, reduce the dosing frequency to every other day.

No dosage modifications are required in other patients with hepatic impairment of any level, or when ETCAMAH is combined with abemaciclib or palbociclib, although patients with moderate or severe hepatic impairment (Child-Pugh B or C) should be monitored for increased adverse reactions and dosage modified as recommended.

Please see Important Safety Information on pages 4-5 and complete Prescribing Information, including Boxed WARNING, and Patient Information.

You may report side effects related to AstraZeneca products .

For more information, please contact AstraZeneca Access 360™ at **1-844-ASK-A360**, Monday through Friday, 8 AM - 6 PM ET.



1-844-ASK-A360 (1-844-275-2360)



1-844-FAX-A360 (1-844-329-2360)



www.MyAccess360.com



Access360@AstraZeneca.com



One MedImmune Way, Gaithersburg, MD 20878

References: **1.** ETCAMAH® (camizestrant) [prescribing information]. Wilmington, DE: AstraZeneca Pharmaceuticals LP; 2026. **2.** Centers for Medicare & Medicaid Services. 2026 ICD-10-CM. Accessed June 1, 2026. <https://www.cms.gov/medicare/coding-billing/icd-10-codes>.



ETCAMAH is a registered trademark and AstraZeneca Access 360 is a trademark of the AstraZeneca group of companies.

©2026 AstraZeneca. All rights reserved. US-102019 Last Updated 9/26